

X-RAY TIMING, SPECTROSCOPY AND PHOTOMETRY OF THE ANOMALOUS X-RAY PULSAR CANDIDATE CXOU J010043.1–721134

M. B. MCGARRY AND B. M. GAENSLER¹, S. M. RANSOM^{2,3} AND V. M. KASPI², AND S. VELJKOVIC⁴

Draft version February 2, 2008

ABSTRACT

We present new X-ray timing and spectral results on the 8.0-second X-ray pulsar CXOU J010043.1–721134 from a series of observations using the *Chandra X-ray Observatory*. We find a spin period in 2004 January of 8.020392 ± 0.000009 seconds. Comparison of this to 2001 *Chandra* observations implies a period derivative $\dot{P} = (1.88 \pm 0.08) \times 10^{-11} \text{ s s}^{-1}$, leading to an inferred dipole surface magnetic field of $3.9 \times 10^{14} \text{ G}$. The spectrum is well fit to an absorbed blackbody of temperature $kT = 0.38 \pm 0.02 \text{ keV}$ with a power law tail of photon index $\Gamma = 2.0 \pm 0.6$. We find that the source has an unabsorbed X-ray flux (0.5–10 keV) of $4_{-1}^{+2} \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ and a corresponding X-ray luminosity of $\sim 2 \times 10^{35} \text{ erg s}^{-1}$ for a distance of 60 kpc. These properties support classification of CXOU J010043.1–721134 as the seventh confirmed anomalous X-ray pulsar, the eleventh confirmed magnetar, and the first magnetar to be identified in the Small Magellanic Cloud.

Subject headings: pulsars: general — pulsars: individual(CXOU J010043.1–721134) — stars: neutron

1. INTRODUCTION

Before the mid-1990s, it was thought that young neutron stars had short ($P \ll 1\text{s}$) spin periods, surface magnetic fields $B \sim 10^{12} \text{ G}$, and exhibited radio pulsations. However, recent theoretical and observational work has revised this picture. We now know that a small population of sources known as soft gamma repeaters (SGRs) are young neutron stars with extreme ($B \sim 10^{14-15} \text{ G}$) surface magnetic fields, or “magnetars”. Another class of objects, anomalous X-ray pulsars (AXPs), are thought to be closely related to SGRs. Anomalous X-ray pulsars are so-named due to their high X-ray luminosities and unusually fast spindown rates, with no evidence of variation due to binary motion, which distinguish them from both isolated radio pulsars and accreting X-ray binaries. Most significantly, AXPs exhibit a small range of spin periods (5–12 seconds) with steady spindown that, if powered by magnetic dipole radiation, imply magnetic fields 2–3 orders of magnitude larger than those of typical radio pulsars (Thompson & Duncan 1996). The magnetar model for AXPs is supported by the SGR-like bursting activity that has been observed from some of these sources (Gavriil, Kaspi, & Woods 2002; Kaspi et al. 2003). Other characteristics include spectra that are well fit by a blackbody model with a temperature $kT \sim 0.5 \text{ keV}$ and a power law tail with photon index $\Gamma=2-4$, and lack of a measurable binary companion that could otherwise produce similar X-ray luminosities through accretion (see Woods & Thompson 2005, for a summary of the properties of known magnetars). With unabsorbed luminosities in the range of 10^{34} to $10^{36} \text{ erg s}^{-1}$, inferred magnetic fields of 10^{14} – 10^{15} Gauss , and characteristic ages of $\sim 8 - 200 \text{ kyr}$, it has been firmly established that AXPs belong to the

magnetar family (Gavriil, Kaspi, & Woods 2004).

The X-ray source CXOU J010043.1–721134 in the Small Magellanic Cloud (SMC) was first detected by *Einstein* in 1979, and was later observed several times by *ROSAT* and *ASCA*, but its periodicity was not noticed until an archival search for X-ray pulsations was carried out on a 2001 *Chandra* observation of a nearby field (Lamb et al. 2002). The period of this source was determined to be 8.0 seconds and its X-ray luminosity was measured to be on the order of $10^{35} \text{ erg s}^{-1}$, assuming a distance to the SMC of 60 kpc (Lamb et al. 2002, 2003; Majid, Lamb, & Macomb 2004). Such a high luminosity and long spin period suggested that this object may belong to the growing class of AXPs (Lamb et al. 2002; Majid, Lamb, & Macomb 2004).

In this Letter we present new *Chandra* results supporting CXOU J010043.1–721134 as the newest member of the AXP class. We present timing analysis from which we measure a period derivative, spectral results, and an analysis of archival *Chandra* and *XMM* data that assesses the flux characteristics of the source over time.

2. OBSERVATIONAL SETUP

Five new *Chandra* observations of CXOU J010043.1–721134 were completed between 2004 January 27 and 2004 March 25. The source was clearly detected at each epoch, located at (J2000) right ascension $01^{\text{h}}00^{\text{m}}43^{\text{s}}.03$, declination $-72^{\circ}11'33''.6$, with a radial uncertainty of $0''.5$. Coordinates from the *Chandra* pointings have not yet been refined using an optical reference. Each observation had approximately 16 ks of on-source time, and the observations were spaced in a geometric series designed for a phase-coherent timing analysis. All five observations were made with the ACIS-S3 CCD in Very Faint mode. A 1/8 subarray was used to reduce the time resolution to 0.4 s.

Additionally, we analyzed serendipitous archival observations from both *Chandra* and *XMM*. The archival 2001 May 15 *Chandra* observation was made in faint mode, and the source fell on the ACIS-I1 CCD approximately $10'$ from the aimpoint. We reprocessed the level 1

¹ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138

² Department of Physics, McGill University, Rutherford Physics Building, 3600 University Street, Montreal, QC H3A 2T8, Canada

³ Current Address: National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA 22903

⁴ Harvard College, Cambridge, MA 02138

events file and applied the CTI, standard gain, and time-dependent gain corrections. *XMM* archival observations were taken on 2000 October 10 and on 2001 November 20 using the EPIC MOS detectors, and the 2001 observation also used the EPIC PN detector. Both datasets lost some observation time due to high background. The source location in each *XMM* observations is also off-axis, and the source lies on a chip gap on the MOS2 instrument. Therefore, we discarded all MOS2 data in this analysis. Table 1 contains detailed information on each observation. We processed *Chandra* observations using CIAO 3.1 and CALDB 2.28, and *XMM* observations using SAS 6.0.0.

3. TIMING ANALYSIS

The five 2004 *Chandra* observations of CXOU J010043.1–721134 were separated by successive intervals of 0.40, 1.72, 8.9, and 46.4 days. For each epoch, we extracted the events from a 2'' radius region centered on the coordinates of the source and corrected their arrival times to the solar system barycenter. We measured pulse profiles for each observation by folding the X-ray events at a range of periods around the $P \sim 8.02$ s spin-period of the pulsar as reported by Lamb et al. (2002, 2003). The profiles with the highest significance (as determined by maximizing χ^2 with respect to a model with no pulsations) were cross-correlated and co-added in phase to create an integrated template pulse profile from all the observations. We then determined pulse arrival times by cross-correlating the profiles from each observation with the high signal-to-noise template profile. From the three *Chandra* observations in 2004 January, we unambiguously measure a spin period $P = 8.020392(9)$ s, where the number in parentheses indicates the uncertainty in the last digit. Unfortunately, the fractional phase errors from these measurements were ~ 0.06 , significantly larger than the 0.02–0.03 fractional phase errors that we were expecting. These larger arrival time uncertainties mean that the spin period determined above from the three 2004 January observations was not of sufficient accuracy to unambiguously account for each rotation of the pulsar between the 2004 January and February observations, let alone during the much longer gap between the 2004 February and March observations.

From the archival *Chandra* observation taken on 2001 May 15, we measure $P = 8.0188(1)$ s, which is significantly shorter than $P = 8.020392(9)$ s in 2004 January as determined above. If this period difference is due to a relatively constant spin-down, it implies $\dot{P} = 1.9(1) \times 10^{-11}$ s s $^{-1}$. If we then force a phasing of the five 2004 observations to most closely match the average spin-down between 2001 May and 2004 January, we determine $\dot{P} = 1.88(8) \times 10^{-11}$ s s $^{-1}$. A single phase wrap between the 2004 February and 2004 March observations implies $\dot{P} = 1.26(8) \times 10^{-11}$ s s $^{-1}$ or $\dot{P} = 2.50(8) \times 10^{-11}$ s s $^{-1}$, both of which are clearly inconsistent with the average spin-down. It is therefore highly likely that the true spin-down value of this source is $\dot{P} = 1.88(8) \times 10^{-11}$ s s $^{-1}$.

4. SPECTROSCOPY

For the spectral analysis on the 2004 *Chandra* observations, we used an extraction radius of 2'' for the source region and 29'03 for the background region. For the

2001 off-axis *Chandra* observation, the source region was an ellipse with major and minor axes 10'82 and 9'84 respectively, and background extraction circle with radius 92'50. In all *Chandra* analyses, the background regions were not centered on the source but did cover the source and so the source region was excluded. For the *XMM* observations, the source extraction circles for the MOS and PN detectors had radii of 12'40 and 19'20, respectively, while the background regions for both were 65'60 and did not overlap the source region.

We successfully fit each spectrum in Sherpa⁵ to a model containing an absorbed blackbody plus a power law, accounting for absorption from the Milky Way and the SMC. We assumed a Galactic absorbing column $N_H = 4.3 \times 10^{20}$ cm $^{-2}$ (Nazé et al. 2003). Standard SMC elemental abundances were used the hydrogen column due to the SMC contribution was allowed to vary⁶ (Russell & Dopita 1992). The temperature and photon index, and their normalizations, were allowed to vary in their respective models.

Table 2 compares various models that we fit to the data. A blackbody plus power law model provides a much better fit than the single model fits that were previously attempted (Lamb et al. 2002; Nazé et al. 2003; Majid, Lamb, & Macomb 2004). Fitting to a simple absorbed power law model yields a poor result, as shown by the reduced $\chi^2 = 2.12$. Fitting to a simple absorbed blackbody model yields a reduced $\chi^2 = 1.10$. However, this fit is infeasible because it requires that there be no absorption column contribution from the SMC (N_{HSMC}). In fact, if N_{HSMC} is frozen at a reasonable value we are still not able to fit a sensible model to the data.

Statistics were calculated to 90% confidence ($\sigma = 1.6$). In each dataset, we excluded data below 0.5 keV to reflect telescope sensitivity and above 4.0 keV because the source spectrum is not detected beyond this. We subtracted the background, although it was minimal in the *Chandra* data, and grouped the spectra into bins of at least 25 counts for analysis.

4.1. Chandra 2004 Observations

We fit the five 2004 *Chandra* datasets simultaneously and find that $N_{HSMC} = (3 \pm 4) \times 10^{21}$ cm $^{-2}$. The large uncertainty in this absorbing column is due to the coupling between the absorbing column and the power law index Γ . We find a blackbody temperature $kT = 0.38 \pm 0.02$ keV, power law photon index $\Gamma = 2.0 \pm 0.6$, and reduced $\chi^2 = 0.97$. The unabsorbed 0.5–10 keV luminosity of the source in these *Chandra* observations is $\sim 2 \times 10^{35}$ erg s $^{-1}$ at a distance of 60 kpc, and the blackbody component contributes $\sim 50\%$ of the luminosity.

4.2. 2000 and 2001 Archival Observation

We also analyzed the 2001 archival *Chandra* observation. The model applied to the previously described

⁵ <http://cxc.harvard.edu/sherpa/>

⁶ Abundances for elements Co and Ca were not included in the model of Russell & Dopita (1992), and so both were assigned a value of 0.3171 with respect to solar in the spectral model based on the average abundance of the other elements. Variations in the Co and Ca abundances do not result in significant changes to the fits.

Chandra data also yields an excellent fit to this observation, with $N_{HSMC} = (8 \pm 6) \times 10^{21} \text{ cm}^{-2}$, $kT = 0.36 \pm 0.03 \text{ keV}$, $\Gamma = 1.9 \pm 0.4$, and reduced $\chi^2 = 1.15$. The unabsorbed 0.5–10 keV luminosity of the source is $\sim 2 \times 10^{35} \text{ erg s}^{-1}$, and the blackbody component contributes $\sim 60\%$ of the luminosity. It should be noted that these data were fit after applying a user-contributed time-dependent gain correction. This software accounts for an instrumental effect that has not been fully addressed by the default Sherpa package for older observations.

The best fitting model to the 2000 and 2001 *XMM* data is again the absorbed blackbody with a power law component. Though we also fit the two *XMM* datasets individually, we found that they were well modeled when fit jointly. This model implies $N_{HSMC} = (5 \pm 7) \times 10^{21} \text{ cm}^{-2}$, $kT = 0.31 \pm 0.04 \text{ keV}$, $\Gamma = 2.0 \pm 0.5$, and reduced $\chi^2 = 1.08$. The unabsorbed 0.5–10 keV luminosity of the source as seen by *XMM* is $\sim 2 \times 10^{35} \text{ erg s}^{-1}$, and the blackbody component contributes $\sim 40\%$ of the luminosity.

5. LONG TERM FLUX BEHAVIOR

We have also examined the flux variability of CXOU J010043.1–721134 over the last three decades using the data in §4, plus count rates from archival data reported by Lamb et al. (2002). We inferred the fluxes and their errors for the older observations using the following method. For both the *Chandra* and *XMM* observations presented in this paper, we used Sherpa’s eflux command to calculate the flux for the best-fit parameters of N_{HSMC} , Γ , and kT . We also determined minimum and maximum possible fluxes, by applying the errors in the spectral fits, which were calculated at the 90% confidence level, in order to achieve the largest possible flux range. Specifically, the minimum flux occurs when N_{HSMC} and Γ are at a maximum and kT is at a minimum, and the maximum flux occurs in the reverse scenario. We applied a correction to account for the correlations in the uncertainties of the three parameters and defined the flux error to be the difference between the flux defined by the best-fit parameters and these extreme flux values.

Fluxes for the observations from previous X-ray satellites listed in Lamb et al. (2002) could not be calculated in this manner without reprocessing each dataset, so we developed a method by which to extrapolate their fluxes and errors. We converted the errors from our *Chandra* observations into fractional uncertainties and then compared them to the fractional uncertainty in their measured count rate to obtain a count rate-flux uncertainty relation. Using the best-fit model for the 2004 *Chandra* observations (see Table 1), we converted from count rate to flux using HEASARC’s WebPIMMS⁷. We then used our count rate-flux relation to extrapolate the uncertainties in these older observations. Results are plotted in Figure 1. Because small changes in the fit parameters lead to large variation in flux for a given count rate, it is difficult to compare fluxes across multiple epochs. However, at face value, the flux behavior does not seem to demonstrate variation spanning more than one order of magnitude over a timescale of ~ 25 years.

6. DISCUSSION

The properties of CXOU J010043.1–721134 are completely consistent with those expected of an AXP. In 2004 January, the source had a period of 8.020392(9) seconds and a $\dot{P} = 1.88(8) \times 10^{-11} \text{ s s}^{-1}$. If this spin-down is due to “standard” dipole radiation, the implied surface magnetic field strength is $3.2 \times 10^{19} (P\dot{P})^{1/2} \text{ G} = 3.9 \times 10^{14} \text{ G}$, consistent with magnetic fields of other AXPs. Using the estimated period derivative, it is possible to calculate the characteristic age and spindown luminosity of the source. A period derivative $\dot{P} = 1.88(8) \times 10^{-11} \text{ s s}^{-1}$ implies a characteristic age of $P/2\dot{P} = 6800$ years (with a 4% uncertainty), consistent with characteristic ages of magnetars (Gaensler et al. 2001). CXOU J010043.1–721134 has a spin-down luminosity of $4\pi^2 I \dot{P} / P^3 \equiv 1 \times 10^{33} \text{ erg s}^{-1}$, where $I \equiv 10^{45} \text{ g cm}^2$ is the assumed moment of inertia of the neutron star.

The source spectrum is well described by a photon index of $\Gamma = 2.0 \pm 0.6$ and a blackbody temperature of $kT = 0.38 \pm 0.02 \text{ keV}$. This implies a 0.5–10 keV unabsorbed luminosity of $\sim 2 \times 10^{35} \text{ erg s}^{-1}$, where the blackbody contributes $\sim 50\%$ of the luminosity (at a distance of 60 kpc). This fit is consistent with those of other AXPs, which tend to have $kT \sim 0.5 \text{ keV}$ and $\Gamma \sim 2$ –4. Although the blackbody contribution to the luminosity of an AXP is highly energy dependent, the value of $\sim 50\%$ in the case of CXOU J010043.1–721134 is consistent with other AXPs in the 1–4 keV range (Özel, Psaltis, & Kaspi 2001).

It has been suggested that the source could be a Be X-ray binary with a ~ 25 day period because there is an optical Be star within $2''$ of the source (Nazé et al. 2003). X-ray binaries are also described by a blackbody plus power law model, but generally their blackbody temperature is higher (i.e. 1–2 keV) (Haberl & Pietsch 2004). However, our timing analysis rules out a 10–30 day binary due to the absence of acceleration by a massive companion in the first three *Chandra* observations since the phase connects unambiguously over ~ 3 days. The phase wrap after the third observation does leave open the possibility of longer period binary motion, however, the long term period change seems to match the best phase connected solution, which likely rules out any binary companion and argues strongly for a relatively steady spin-down. We further note that Durant & van Kerkwijk (2005) have identified a likely optical counterpart to CXOU J010043.1–721134 distinct from the adjacent B star. This object provides a convincing alternative to the Be X-ray binary interpretation.

CXOU J010043.1–721134 does not show strong evidence of variability over the timescale of years. Although not all AXPs are variable, some AXPs have been found to vary across one to two orders of magnitude (Gavril, Kaspi, & Woods 2004; Ibrahim et al. 2004). X-ray binaries also exhibit variations, by factors of up to 20 over a similar timescale, and can change by a factor of more than 100 during giant outbursts (Haberl & Pietsch 2004). Even considering the large uncertainties, we do not see variability on this scale in CXOU J010043.1–721134 over the past 25 years. While it does not conclusively characterize the object, apparent lack of variability across orders of magnitude does not challenge the categorization of this source as an AXP.

In summary, the spectral, timing, and photometric properties of CXOU J010043.1–721134 are consistent

⁷ <http://heasarc.gsfc.nasa.gov/Tools/w3pimms.html>

TABLE 1
SPECTRAL FITS TO *Chandra* AND *XMM* DATASETS

Dataset & Obsid	Date	On-Source Time (ks)	Count Rate (10^{-2} ct s $^{-1}$)	N_{HSMC} (10^{21} cm $^{-2}$)	Γ	kT_{BB} (keV)	χ^2/DOF
XMM 0110000201	2000 Oct 10	14.5	5.4(2)	1(5)	1.8(1)	0.35(6)	21.2/21
CXO 1881	2001 May 15	98.7	6.03(8)	8(6)	1.9(4)	0.36(3)	130/113
XMM 0018540101 ^a	2001 Nov 20	24.7/20.2	4.0(1)/15.1(3)	6(8)	2.0(3)	0.30(5)	131/126
CXO 4616	2004 Jan 27	16.2	8.8(5)	3(4)	2.0(6)	0.38(2)	45.9/38
CXO 4617	2004 Jan 28	16.1	8.8(5)	3(4)	2.0(6)	0.38(2)	40.8/39
CXO 4618	2004 Jan 29	16.5	8.5(5)	3(4)	2.0(6)	0.38(2)	34.3/39
CXO 4619	2004 Feb 07	16.7	8.6(5)	3(4)	2.0(6)	0.38(2)	35.8/39
CXO 4620	2004 Mar 25	15.6	9.2(5)	3(4)	2.0(6)	0.38(2)	49.0/37

NOTE. — Parentheses following each value represent the error in the last digit; “On-Source Time” refers to the amount of usable time recovered from the original observation, excluding deadtime caused by CCD readout, high background periods, etc.; Count Rate is in the default energy range for each instrument (*Chandra*: 0.2–10keV, *XMM*: 0.4–10keV); N_{HSMC} = the absorbing column due to the Small Magellanic Cloud only, assuming a Galactic absorbing column of 4.3×10^{20} cm $^{-2}$; Γ = the power law photon index; kT_{BB} = blackbody characteristic temperature; χ^2 value is for the fit of the absorbed black body plus power law model; The unabsorbed 0.5–10keV flux for the five 2004 *Chandra* observations is $4^{+2}_{-1} \times 10^{-13}$ erg cm $^{-2}$ s $^{-1}$, and in all other observations is $5^{+2}_{-1} \times 10^{-13}$ erg cm $^{-2}$ s $^{-1}$.

^aValues for MOS1 and PN respectively.

with the properties of other AXPs (Mereghetti et al. 2002), expanding this class of magnetars to seven confirmed members. We note that the source is the only known magnetar in the SMC. If the magnetar birth rate is $\sim 10\%$ of that of radio pulsars (Kouveliotou et al. 1994; Gaensler et al. 2005), then for an SMC supernova rate of $(5 \pm 3) \times 10^{-4}$ per year (Crawford et al. 2001) and a magnetar lifetime on the order of 10^4 yr (Colpi, Geppert, & Page 2000; Gaensler et al. 2001), we can expect 0.5 ± 0.3 magnetars to exist in the SMC at a given time. This would be consistent with the fact that only this magnetar is known in the SMC thus far, despite several archival searches for X-ray sources (Majid, Lamb, & Macomb 2004; Haberl & Pietsch 2004; Laycock et al. 2005). Interestingly, (Ibrahim et al. 2004) report the discovery of a transient magnetar, detected

only because of an outburst that resulted in a substantial flux enhancement that faded on a time scale of months. Similarly, the highly variable 7-s pulsar AX J1845-0258 may be another such source (Vasisht et al. 2000). These sources suggest that the magnetar birthrate could be substantially higher than has been estimated, possibly even comparable to the radio pulsar birthrate. Continued monitoring of the SMC will provide a useful test of these possibilities. In any case, CXOU J010043.1–721134, with a well-known distance and low extinction, will be a valuable probe into this exotic class of objects.

We thank Fernando Camilo, Cara Rakowski, Michael Garcia, and Terry Gaetz for useful discussions, and the *Chandra* Helpdesk for software support. This work was supported by NASA through contract NAS 8-39073 and SAO grant GO4-5065X.

REFERENCES

- Colpi, M., Geppert, U., Page, D. 2000, ApJ, 529, 1, L29
Crawford, F., Kaspi, V. M., Manchester, R. N., Lyne, A. G., Camilo, F., & D’Amico, N. 2001, ApJ, 553, 367
Durant, M. & van Kerkwijk, M. H. 2005, ApJ, submitted
Gaensler, B. M., McClure-Griffiths, N. M., Oey, M. S., Haverkorn, M., Dickey, J. M., & Green, A. J. 2005, ApJ, 620, L95
Gaensler, B. M., Slane, P. O., Gotthelf, E. V., & Vasisht, G. 2001, ApJ, 559, 963
Gavriil, F. P., Kaspi, V. M., & Woods, P. M. 2004, Advances in Space Research, 33, 4, 654
Gavriil, F. P., Kaspi, V. M., & Woods, P. M. 2002, Nature, 419, 142
Haberl, F., & Pietsch, W. 2004, A&A, 414, 667
Ibrahim, A. I., et al. 2004, ApJ, 609, L21
Kaspi, V. M., Gavriil, F. P., Woods, P. M., Jensen J. B., Roberts, M. S. E., & Chakrabarty, D. 2003, ApJ, 588, L93
Kouveliotou, C., et al 1994, Nature, 368, 125
Lamb, R. C., Fox, D. W., Macomb, D. J., & Prince, T. A. 2002, ApJ, 574, L29
— 2003, ApJ, 599, 2, L115
Laycock, S., Corbet, R. H. D., Coe, M. J., Marshall, F. E., Markwardt, C., & Lochner, J. 2005, ApJ, in press, (astro-ph/0406420)
Majid, W. A., Lamb, R. C., & Macomb, D. J. 2004, ApJ, 609, 133
Mereghetti, S., Chialone, L., Israel, G. L., & Stella, L. 2002, in Conf. Ser. 270, WE-Heraeus Seminar on Neutron Stars, Pulsars, and Supernova Remnants, MPE Report 278, ed. W. Becker, J. Lesch, & J. Trümper (Munich: Max-Planck-Institute), 29
Nazé, Y., Hartwell, J. M., Stevens, I. R., Manfroid, J., Marchenko, S., Corcoran, M. F., Moffat, A. F. J., & Skalkowski, G. 2003, ApJ, 586, 983
Özel, F., Psaltis, D., & Kaspi, V. M. 2001, ApJ, 563, 255
Russell, S. C., & Dopita, M. A. 1992, ApJ, 384, 508
Thompson, C., & Duncan, R. C. 1996, ApJ, 473, 322
Vasisht, G., Gotthelf, E. V., Torri, K., & Gaensler, B. M. 2000, ApJ, 542, L49
Woods, P. M., & Thompson, C. 2004, to appear in “Compact Stellar X-ray Sources”, eds. W.H.G. Lewin and M. van der Klis, in press, (astro-ph/0406133)

TABLE 2
JOINT SPECTRAL FITS TO 2004 *Chandra* DATA

Model	$N_{HSMC}(\text{cm}^{-2})$	$\Gamma/kT_{BB}(\text{keV})$	F ($10^{-13} \text{ erg cm}^2 \text{ s}^{-1}$)	χ^2/DOF
BB	... ^a	... /0.40(8)	~ 4	235/214 = 1.1
PL	$8(1) \times 10^{21}$	2.20(5)/...	~ 6	455/214 = 2.1
BB+PL	$3(4) \times 10^{21}$	2.0(6)/0.38(2)	~ 4	205/212 = 0.97

NOTE. — Parentheses following each value represent the error in the last digit; N_H = the absorbing column due to the Small Magellanic Cloud only, assuming a Galactic absorbing column of $4.3 \times 10^{20} \text{ cm}^{-2}$; Γ = the power law photon index; kT_{BB} = blackbody characteristic temperature; F = Unabsorbed flux for the model in the 0.5–10 keV range.

^aThe data could not be fit within Sherpa’s limits for N_{HSMC} .




FIG. 1.— Estimated unabsorbed fluxes for CXOU J010043.1–721134 over the last three decades of observations, using the BB+PL model from Table 2 in the energy range 0.2–10 keV. Each instrument is denoted with a separate plot symbol as follows: X - *Einstein* IPC, $\diamond$ - *ROSAT* PSPC, $\triangle$ - *ROSAT* HRI, $\square$ - *ASCA* GIS, * - *XMM* MOS/PN, $\bullet$ - *Chandra* ACIS. Flux estimates from observations presented in this *Letter* were determined by using Sherpa’s eflux command. Those not presented in this *Letter* are derived from the observed count rates listed in Lamb et al. (2002) by using HEASARC WebPIMMS.